

Dalton Communications

Synthesis of the First Organometallic Monosubstituted Octanuclear Silasesquioxane

Gion Calzaferri* and Roman Imhof

Institute of Inorganic and Physical Chemistry, University of Berne, CH-3000 Berne 9, Switzerland



The hexachloroplatinic acid-catalysed hydrosilylation of vinylferrocene by octahydrosilasesquioxane $H_8Si_8O_{12}$ to the first organometallic monosubstituted octasilasesquioxane $[(\eta-C_5H_5)Fe(\eta-C_5H_4CH_2CH_2)]H_7Si_8O_{12}$ has been achieved.

The structure of the highly symmetrical octahydrosilasesquioxane $H_8Si_8O_{12}$ was first determined by Larsson^{1a} and has lately been reinvestigated.^{1b} It is one of the most appealing members of the class of silasesquioxanes of the type $(RSiO_3/2)_n$, $n = 1, 2, 3, \text{ etc.}$, and can be prepared in good yields by the polycondensation of $SiHCl_3$ in a biphasic solution.² This class of substances³ seemed to be forgotten for a long time. Structurally well defined silasesquioxanes and silicate clusters have, however, recently attracted interest as 'building blocks' for the preparation of highly siliceous materials.⁴ Agaskar⁵ has developed an elegant simple procedure for the synthesis of a wide variety of silylated spherosilicates, many of which offer interesting possibilities as precursors to organolithic macromolecular material. New organosiliceous polymeric materials have been prepared by Hoebbel *et al.*⁶ Feher *et al.*^{7a} followed a different route by using an incompletely condensed heptamer trisilanol as a model for a silicon surface on which they examined metal catalysed reactions. These investigations led to the synthesis of some new metallasilasesquioxanes with important implications for the chemistry of silica surfaces.^{7b}

Our interest for the silasesquioxanes has been attracted by the resemblance of $H_8Si_8O_{12}$ to the double-four ring of the secondary building unit of zeolite A and by the appealing polycyclic structure which makes these compounds attractive as models for vibrational spectroscopy.⁸ Furthermore, the possible use of $H_8Si_8O_{12}$ as a framework for building donor-acceptor systems encouraged us to explore the chemistry of this substance. After the palladium-catalysed deuterium exchange and the substitution of all H atoms by alkyl groups, we recently found a way to prepare $RH_7Si_8O_{12}$.⁹ It became clear that a great number of interesting compounds would become accessible if a route to organometallic monosubstituted octasilasesquioxanes could be found. Two main ways seemed to be within reach. One was to start with $RH_7Si_8O_{12}$, e.g. $R = (CH_2)_2Ph$, and to treat it with e.g. $[Cr(CO)_6]$. The other possibility was to try hydrosilylation by treating $H_8Si_8O_{12}$ with a vinyl-metalloorganic compound as shown in Scheme 1. To test the second reaction path, we tried a hexachloroplatinic acid-

catalysed hydrosilylation of the well understood vinylferrocene by $H_8Si_8O_{12}$ and found that an easy one-flask reaction leads to the desired monosubstituted organometallic octasilasesquioxane $[(\eta-C_5H_5)Fe(\eta-C_5H_4CH_2CH_2)]H_7Si_8O_{12}$ **1**.†

We assume that the mechanism of this reaction is dictated by the rigidity and cage size of $H_8Si_8O_{12}$ and that it proceeds *via* five-co-ordinated Si similar to the reaction path for the palladium-catalysed deuterium exchange of $H_8Si_8O_{12}$ to $D_8Si_8O_{12}$.¹¹ The structure of compound **1** is unequivocally demonstrated by ¹H, ¹³C and ²⁹Si NMR and mass spectroscopy. The ¹H NMR spectrum shows the expected splitting pattern, especially two singlets for the SiH protons with relative intensities of 4:3. The CH_2CH_2 bridge between the octa-



Scheme 1 Hydrosilylation, a reaction path to monosubstituted organometallic silasesquioxanes; e.g. $R = \text{ferrocenyl}$

† The compound $H_8Si_8O_{12}$ (100 mg, 0.24 mmol)² was dissolved in a round-bottom flask (25 cm³) in toluene (15 cm³), vinylferrocene (50 mg, 0.24 mmol) (Aldrich) and H_2PtCl_6 solution (10 μ l, 0.01 mol dm⁻³ in Pr^iOH) were added in toluene (4 cm³) and the mixture heated at reflux for 20 h under N_2 . After evaporation of the solvent a yellow solid precipitated which was suspended in hexanes (Romil Chemicals), filtered and the remaining yellow solution injected on a 600 \times 25 mm PolymerLab (pore size 50 \AA , particle size 10 μ m) size-exclusion liquid chromatography column. The yield of compound **1** was 21 mg (0.035 mmol, 14%). IR: see lit.,¹⁰ NMR ($CDCl_3$, standard $SiMe_4$: ¹H (300 MHz), δ 1.01 (m, 2 H, $SiCH_2$), 2.46 (m, 2 H, $CH_2C_5H_4$), 4.06 (m, 4 H, $CH_2C_5H_4$), 4.11 (s, 5 H, C_5H_5), 4.23 (s, 3 H, SiH) and 4.25 (s, 4 H, SiH); ¹³C (75 MHz), δ 13.0 (1 C, $SiCH_2$), 22.3 (1 C, $CH_2C_5H_4$), 67.3, 67.8 (4 C, $CH_2C_5H_4$), 68.5 (5 C, C_5H_5) and 90.8 (1 C, $CH_2C_5H_4$); ²⁹Si (60 MHz, inverse gated, external standard), δ -66.7 (1 Si, $SiCH_2$), -84.58 (3 Si), -84.60 (3 Si) and -84.64 (1 Si). Solutions of **1** in chlorinated solvents should be kept in the dark. Mass spectrum (70 eV, ca. 1.12×10^{-17} J); m/z 636 (100, M^+), 423 (16, $[M - C_{12}H_{13}Fe]^+$), 199 (20, $[C_{11}H_{11}Fe]^+$) and 121 (27%, $[C_5H_5Fe]^+$).

silasesquioxane and the ferrocene forms an AA'XX' system which is responsible for the multiplets in the corresponding regions. The ^{29}Si NMR spectrum shows four peaks with relative intensities of 1:3:3:1. The three Si atoms in the direct neighbourhood of the substituted Si atoms have the largest shift with respect to unsubstituted $\text{H}_8\text{Si}_8\text{O}_{12}$. The IR spectrum in the range 350–4000 cm^{-1} shows the expected pattern to be discussed later.¹⁰

Acknowledgements

We thank Dr. Ch. Müller for measuring the ^{29}Si NMR spectrum and for helpful discussions of the NMR results. This work is part of project NF 20-28528.90, financed by the Schweizerischer Nationalfonds zur Förderung der wissenschaftlichen Forschung and project BEW-EPA 217.307, financed by the Schweizerisches Bundesamt für Energiewirtschaft.

References

- (a) K. Larsson, *Ark. Kemi*, 1960, **16**, 215; (b) T. P. E. Auf der Heide, H.-B. Bürgi, H. Bürgy and K. W. Törnroos, *Chimia*, 1991, **45**, 38.
- P. A. Agaskar, *Inorg. Chem.*, 1991, **30**, 2702.
- D. W. Scott, *J. Am. Chem. Soc.*, 1946, **68**, 356.
- P. A. Agaskar, V. W. Day and W. G. Klemperer, *J. Am. Chem. Soc.*, 1987, **109**, 5554; V. W. Day, W. G. Klemperer, V. V. Mainz and D. Millar, *J. Am. Chem. Soc.*, 1985, **107**, 8262.
- P. A. Agaskar, *Inorg. Chem.*, 1990, **29**, 1603; *J. Am. Chem. Soc.*, 1989, **111**, 6858.
- D. Hoebbel, I. Pitsch and D. Heidemann, *Z. Anorg. Allg. Chem.*, 1991, **592**, 207.
- (a) F. J. Feher, J. F. Walzer and R. L. Blanski, *J. Am. Chem. Soc.*, 1991, **113**, 3618; (b) F. J. Feher, T. A. Budzichowski, K. Rahimian and J. W. Ziller, *J. Am. Chem. Soc.*, 1992, **114**, 3859.
- P. Bornhauser and G. Calzaferri, *Spectrochim. Acta, Part A*, 1990, **46**, 1045; M. Bärtsch, P. Bornhauser, H. Bürgy and G. Calzaferri, *Spectrochim. Acta, Part A*, 1991, **47**, 1627; M. Bärtsch, P. Bornhauser and G. Calzaferri, *SPIE J.*, 1992, **1575**, 588.
- H. Bürgy and G. Calzaferri, *Helv. Chim. Acta*, 1990, **73**, 698; D. Herren, H. Bürgy and G. Calzaferri, *Helv. Chim. Acta*, 1991, **74**, 24; G. Calzaferri, D. Herren and R. Imhof, *Helv. Chim. Acta*, 1991, **74**, 1278.
- M. Bärtsch, P. Bornhauser, G. Calzaferri and R. Imhof, unpublished work.
- G. Calzaferri and R. Hoffmann, *J. Chem. Soc., Dalton Trans.*, 1991, 917.

Received 1st July 1992; Communication 2/03489A